

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark
Office
(Box PCT)
Crystal Plaza 2
Washington, DC 20231
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

Date of mailing: 29 July 1999 (29.07.99)	
International application No.: PCT/EP99/00155	Applicant's or agent's file reference: PRAFF 26/WO
International filing date: 13 January 1999 (13.01.99)	Priority date: 21 January 1998 (21.01.98)
Applicant: SMITS, Georges et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International preliminary Examining Authority on:
29 June 1999 (29.06.99)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer: J. Zahra Telephone No.: (41-22) 338.83.38
---	---

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

PCT

REQUEST

The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty.

For receiving Office use only

International Application No.

International Filing Date

Name of receiving Office and "PCT International Application"

Applicant's or agent's file reference
(if desired) (12 characters maximum) PRAFF 26/W0

Box No. I TITLE OF INVENTION

Process for the manufacture of chicory inulin, hydrolysates and derivatives of inulin, and improved chicory inulin products, hydrolysates and derivatives.

Box No. II APPLICANT

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

TIENSE SUIKERRAFFINADERIJ NV
Tervurenlaan 182
B-1150 Brussel
Belgium

☐ This person is also inventor.

Telephone No.
32-(0)2-775.80.20

Facsimile No.

Teleprinter No.

State (that is, country) of nationality:

BE

State (that is, country) of residence:

BE

This person is applicant for the purposes of:

☐ all designated States

☒ all designated States except the United States of America

☐ the United States of America only

☐ the States indicated in the Supplemental Box

Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

SMITS Georges
Dr. De Cockstraat 16
B-9308 Gijzegem (Aalst), Belgium

This person is:

☐ applicant only

☒ applicant and inventor

☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:

BE

State (that is, country) of residence:

BE

This person is applicant for the purposes of:

☐ all designated States

☐ all designated States except the United States of America

☒ the United States of America only

☐ the States indicated in the Supplemental Box

☒ Further applicants and/or (further) inventors are indicated on a continuation sheet.

Box No. IV AGENT OR COMMON REPRESENTATIVE; OR ADDRESS FOR CORRESPONDENCE

The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:

☒ agent

☐ common representative

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country.)

HERMANS Johny
Tiense Suikerraffinaderij N.V.
Aandorenstraat 1
B-3300 TIENEN
Belgium

Telephone No.
32(0)16 -80.12.92

Facsimile No.
32(0)16 -80.13.59

Teleprinter No.

☐ Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.



Continuation of Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

If none of the following sub-boxes is used, this sheet should not be included in the request.

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

DE LEENHEER Leen
Jezus Eiklaan 64
B-3080 Tervuren, Belgium

This person is:

- ☐ applicant only
☒ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality: BE

State (that is, country) of residence: BE

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

This person is:

- ☐ applicant only
☐ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:

State (that is, country) of residence:

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☐ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

This person is:

- ☐ applicant only
☐ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:

State (that is, country) of residence:

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☐ the United States of America only ☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

This person is:

- ☐ applicant only
☐ applicant and inventor
☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:

State (that is, country) of residence:

This person is applicant for the purposes of:

- ☐ all designated States ☐ all designated States except the United States of America ☐ the United States of America only ☐ the States indicated in the Supplemental Box

☐ Further applicants and/or (further) inventors are indicated on another continuation sheet.



Box No.V DESIGNATION OF STATES

The following designations are hereby made under Rule 4.9(a) (mark the applicable check-boxes; at least one must be marked):

Regional Patent

- ☐ **AP ARIPO Patent:** GH Ghana, GM Gambia, KE Kenya, LS Lesotho, MW Malawi, SD Sudan, SZ Swaziland, UG Uganda, ZW Zimbabwe, and any other State which is a Contracting State of the Harare Protocol and of the PCT
- ☐ **EA Eurasian Patent:** AM Armenia, AZ Azerbaijan, BY Belarus, KG Kyrgyzstan, KZ Kazakhstan, MD Republic of Moldova, RU Russian Federation, TJ Tajikistan, TM Turkmenistan, and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT
- ☒ **EP European Patent:** AT Austria, BE Belgium, CH and LI Switzerland and Liechtenstein, CY Cyprus, DE Germany, DK Denmark, ES Spain, FI Finland, FR France, GB United Kingdom, GR Greece, IE Ireland, IT Italy, LU Luxembourg, MC Monaco, NL Netherlands, PT Portugal, SE Sweden, and any other State which is a Contracting State of the European Patent Convention and of the PCT
- ☐ **OA OAPI Patent:** BF Burkina Faso, BJ Benin, CF Central African Republic, CG Congo, CI Côte d'Ivoire, CM Cameroon, GA Gabon, GN Guinea, ML Mali, MR Mauritania, NE Niger, SN Senegal, TD Chad, TG Togo, and any other State which is a member State of OAPI and a Contracting State of the PCT (if other kind of protection or treatment desired, specify on dotted line)

National Patent (if other kind of protection or treatment desired, specify on dotted line):

- | | |
|---|---|
| <input type="checkbox"/> AL Albania | <input type="checkbox"/> LS Lesotho |
| <input type="checkbox"/> AM Armenia | <input type="checkbox"/> LT Lithuania |
| <input type="checkbox"/> AT Austria | <input type="checkbox"/> LU Luxembourg |
| <input checked="" type="checkbox"/> AU Australia | <input type="checkbox"/> LV Latvia |
| <input type="checkbox"/> AZ Azerbaijan | <input type="checkbox"/> MD Republic of Moldova |
| <input type="checkbox"/> BA Bosnia and Herzegovina | <input type="checkbox"/> MG Madagascar |
| <input type="checkbox"/> BB Barbados | <input type="checkbox"/> MK The former Yugoslav Republic of Macedonia |
| <input type="checkbox"/> BG Bulgaria | <input type="checkbox"/> MN Mongolia |
| <input type="checkbox"/> BR Brazil | <input type="checkbox"/> MW Malawi |
| <input type="checkbox"/> BY Belarus | <input type="checkbox"/> MX Mexico |
| <input type="checkbox"/> CA Canada | <input type="checkbox"/> NO Norway |
| <input type="checkbox"/> CH and LI Switzerland and Liechtenstein | <input type="checkbox"/> NZ New Zealand |
| <input checked="" type="checkbox"/> CN China | <input type="checkbox"/> PL Poland |
| <input type="checkbox"/> CU Cuba | <input type="checkbox"/> PT Portugal |
| <input type="checkbox"/> CZ Czech Republic | <input type="checkbox"/> RO Romania |
| <input type="checkbox"/> DE Germany | <input type="checkbox"/> RU Russian Federation |
| <input type="checkbox"/> DK Denmark | <input type="checkbox"/> SD Sudan |
| <input type="checkbox"/> EE Estonia | <input type="checkbox"/> SE Sweden |
| <input type="checkbox"/> ES Spain | <input type="checkbox"/> SG Singapore |
| <input type="checkbox"/> FI Finland | <input type="checkbox"/> SI Slovenia |
| <input type="checkbox"/> GB United Kingdom | <input type="checkbox"/> SK Slovakia |
| <input type="checkbox"/> GE Georgia | <input type="checkbox"/> SL Sierra Leone |
| <input type="checkbox"/> GH Ghana | <input type="checkbox"/> TJ Tajikistan |
| <input type="checkbox"/> GM Gambia | <input type="checkbox"/> TM Turkmenistan |
| <input type="checkbox"/> GW Guinea-Bissau | <input checked="" type="checkbox"/> TR Turkey |
| <input type="checkbox"/> HR Croatia | <input type="checkbox"/> TT Trinidad and Tobago |
| <input type="checkbox"/> HU Hungary | <input type="checkbox"/> UA Ukraine |
| <input type="checkbox"/> ID Indonesia | <input type="checkbox"/> UG Uganda |
| <input type="checkbox"/> IL Israel | <input checked="" type="checkbox"/> US United States of America |
| <input type="checkbox"/> IS Iceland | <input type="checkbox"/> UZ Uzbekistan |
| <input type="checkbox"/> JP Japan | <input type="checkbox"/> VN Viet Nam |
| <input type="checkbox"/> KE Kenya | <input type="checkbox"/> YU Yugoslavia |
| <input type="checkbox"/> KG Kyrgyzstan | <input type="checkbox"/> ZW Zimbabwe |
| <input type="checkbox"/> KP Democratic People's Republic of Korea | |
| <input type="checkbox"/> KR Republic of Korea | |
| <input type="checkbox"/> KZ Kazakhstan | |
| <input type="checkbox"/> LC Saint Lucia | |
| <input type="checkbox"/> LK Sri Lanka | |
| <input type="checkbox"/> LR Liberia | |

Check-boxes reserved for designating States (for the purposes of a national patent) which have become party to the PCT after issuance of this sheet:

☒ IN India

Precautionary Designation Statement: In addition to the designations made above, the applicant also makes under Rule 4.9(b) all other designations which would be permitted under the PCT except any designation(s) indicated in the Supplemental Box as being excluded from the scope of this statement. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit. (Confirmation of a designation consists of the filing of a notice specifying that designation and the payment of the designation and confirmation fees. Confirmation must reach the receiving Office within the 15-month time limit.)



Box No. VI PRIORITY CLAIM		<input type="checkbox"/> Further priority claims are indicated in the Supplemental Box.		
Filing date of earlier application (day/month/year)	Number of earlier application	Where earlier application is:		
		national application: country	regional application: regional Office	international application: receiving Office
item (1) 21 January 1998 (21.01.1998)	98870012.6		EP	
item (2)				
item (3)				

☒ The receiving Office is requested to prepare and transmit to the International Bureau a certified copy of the earlier application(s) (only if the earlier application was filed with the Office which for the purposes of the present international application is the receiving Office) identified above as item(s) 4) EP 98870012.6

* Where the earlier application is an ARIPO application, it is mandatory to indicate in the Supplemental Box at least one country party to the Paris Convention for the Protection of Industrial Property for which that earlier application was filed (Rule 4.10(b)(ii)). See Supplemental Box.

Box No. VII INTERNATIONAL SEARCHING AUTHORITY			
Choice of International Searching Authority (ISA) <small>(if two or more International Searching Authorities are competent to carry out the international search, indicate the Authority chosen; the two-letter code may be used):</small>		Request to use results of earlier search: reference to that search (if an earlier search has been carried out by or requested from the International Searching Authority):	
ISA / EPO		Date (day/month/year)	Number
		25.06.1998	98870012.6
			Country (or regional Office) EP

Box No. VIII CHECK LIST; LANGUAGE OF FILING	
This international application contains the following number of sheets: request : 4 description (excluding sequence listing part) : 36 claims : 7 abstract : 1 drawings : 0 sequence listing part of description : Total number of sheets : 48	This international application is accompanied by the item(s) marked below: 1. <input checked="" type="checkbox"/> fee calculation sheet 2. <input checked="" type="checkbox"/> separate signed power of attorney 3. <input checked="" type="checkbox"/> copy of general power of attorney; reference number, if any: <u>38099</u> 4. <input type="checkbox"/> statement explaining lack of signature 5. <input type="checkbox"/> priority document(s) identified in Box No. VI as item(s): 6. <input type="checkbox"/> translation of international application into (language): 7. <input type="checkbox"/> separate indications concerning deposited microorganism or other biological material 8. <input type="checkbox"/> nucleotide and/or amino acid sequence listing in computer readable form 9. <input checked="" type="checkbox"/> other (specify): <u>European Search Report EP98870012.6</u>
Figure of the drawings which should accompany the abstract:	Language of filing of the international application: <u>English</u>

Box No. IX SIGNATURE OF APPLICANT OR AGENT	
<small>Next to each signature, indicate the name of the person signing and the capacity in which the person signs (if such capacity is not obvious from reading the request).</small>	
 HERMANS Johny Agent; European Patent Attorney No 85220	

For receiving Office use only	
1. Date of actual receipt of the purported international application:	2. Drawings: <input type="checkbox"/> received: <input type="checkbox"/> not received:
3. Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application:	
4. Date of timely receipt of the required corrections under PCT Article 11(2):	
5. International Searching Authority (if two or more are competent): <u>ISA /</u>	6. <input type="checkbox"/> Transmittal of search copy delayed until search fee is paid.

For International Bureau use only
Date of receipt of the record copy by the International Bureau:



PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference PRAFF 26/WO	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 99/ 00155	International filing date (day/month/year) 13/01/1999	(Earliest) Priority Date (day/month/year) 21/01/1998
Applicant TIENSE SUIKERRAFFINADERIJ NV et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.



as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/00155

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C08B37/18 A01G1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A01G C08B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MICHAEL ERNST ET AL.: "Carbohydrate changes in chicory during growth and storage." SCIENTIA HORTICULTURAE, vol. 63, 1995, pages 251-261, XP002069326 see page 252, paragraph 2.1 see page 259 --- -/--	1-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

28 May 1999

Date of mailing of the international search report

07/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lensen, H



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/L 99/00155

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 127, no. 18, 3 November 1997 Columbus, Ohio, US; abstract no. 245507, BAERT JOOST R.A.: "The effect of sowing and harvest date and cultivar on inulin yield and composition of chicory roots." XP002069328 see abstract & IND. CROPS PROD., vol. 6, no. 3,4, 1997, pages 195-199, ---	1-28
Y	WIM VAN DEN ENDE ET AL.: "Fructan synthesizing and degrading activities in chicory roots (Cichorium intybus L.) during field-growth, storage and forcing." PLANT PHYSIOL., vol. 149, 1996, pages 43-50, XP002069327 STUTTGART see summary ---	1-28
Y	EP 0 787 745 A (MCNEIL-PPC) 6 August 1997 see page 9, line 15 - line 32 see page 2, line 38 - line 45 ---	1-28
A	LEEN DE LEENHEER ET AL.: "Progress in the elucidation of the composition of chicory inulin." STARCH/ DIE STÄRKE, vol. 46, no. 5, 1 May 1994, pages 193-196, XP000442587 WEINHEIM, DE -----	22-28



INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/00155

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 787745	A	06-08-1997	
		AU 1249597 A	14-08-1997
		BR 9700887 A	27-10-1998
		CA 2196739 A	06-08-1997
		HU 9700359 A	30-03-1998
		JP 9324002 A	16-12-1997
		NO 970490 A	06-08-1997
		NZ 314187 A	28-01-1999



PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference PRAFF 26/WO	<div style="display: flex; justify-content: space-between;"> <div>FOR FURTHER ACTION</div> <div>See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)</div> </div>	
International application No. PCT/EP99/00155	International filing date (day/month/year) 13/01/1999	Priority date (day/month/year) 21/01/1998
International Patent Classification (IPC) or national classification and IPC C08B37/18		
Applicant TIENSE SUIKERRAFFINADERIJ NV et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 6 sheets, including this cover sheet.

☒ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 6 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 29/06/1999	Date of completion of this report <div style="text-align: center; font-size: 1.2em;">21.03.00</div>
Name and mailing address of the international preliminary examining authority: <div style="display: flex; align-items: center;"> <div> European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465 </div> </div>	Authorized officer Lucchesi-Palli, C Telephone No. +49 89 2399 2093





**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP99/00155

I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

Description, pages:

1-36 as originally filed

Claims, No.:

1-28 as received on 12/02/2000 with letter of 08/02/2000

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims 1 - 28
	No:	Claims
Inventive step (IS)	Yes:	Claims
	No	Claims 1 - 28
Industrial applicability (IA)	Yes:	Claims 1 - 28
	No	Claims



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP99/00155

2. Citations and explanations

see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

see separate sheet



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP99/00155

ad Section V:

Novelty:

The closest prior art is disclosed in the description of the present application on page 8, last paragraph, in which the conventional cultivation of chicory in the northern parts of Europe is disclosed. The claimed seeding times, cultivation time and thus the harvesting time falls within the periods disclosed in the present description. In this paragraph, lines 32 to 34 it is also stated that the end of the growing season is the time when the biomass of the roots ceases to increase significantly, i.e. after about 180 to 200~~0~~ days of growing. Thus it is known from the prior art that the chicory roots can be grown during a period of at least 150 days (including the possibility of growing of at least 180 days). The prior art does consequently not disclose that the weather conditions have to be such that during a period of at least from the beginning of the third month of the growing period till the end of the processing of the roots the FEH gene has not been triggered by the occurrence of low temperatures.

Claims 1 to 28 fulfill the requirements of article 33(2) PCT.

Inventive step:

According to the description of the present application, page 11, first paragraph, it has been found that the degradation of inulin is mainly catalysed by one or more enzymes with exohydrolase activity coded by one or more FEH genes in the chicory genome.

In document D1, Wim van Ende et al.: "Fructan synthesizing and degrading activities in chicory roots (*Cichorium intybus* L.) during field growth, storage and forcing", Plant Physiology, vol 149, 1996 pages 43 - 50, XP002069327, it is stated that the activity of fructan exohydrolase increased rapidly after mid-october (it has to be emphasized that in this period low temperature conditions can be present in mid-Belgium, where the investigation was made; the conditions varying clearly from year to year) and that during cold storage a further rapid increase was detected. In order to avoid degradation of inulin,



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP99/00155

while achieving the optimal inulin harvest, it is evident for the man skilled in the art to harvest and store the chicory roots at the optimal stage which he will find out by normal routine tests depending on the used cultivar and the climatic conditions in which the roots are grown without the exercise of an inventive skill. That any enzyme activity is coded by one or more genes is known to the man skilled in the art. Although the scientific explanation that the FEH gene should not be triggered is not explicitly mentioned in D1, the influence of low temperatures on inulin degradation has been found in D1 and consequently the man skilled in the art will evidently choose the harvesting time in such a way that low temperatures will not degrade inulin to arrive to the maximal possible result. Thus even the fact that it was not known which mechanism was responsible for the inulin degradation, the skilled person will harvest the roots at their optimal yield, which he has to find out by routine analysis of the roots.

Claims 1, 2, 4, 6, 7, 8 do thus not meet the requirements of article 33(3) PCT.

Claims 3 and 5 do not add any inventive features to the claims from which they depend since the DP and the mean DP is depending not only on the cultivation conditions but also on the used cultivar. Since the cultivars used in the present application are known cultivars no surprising effect is to be expected when cultivating the known cultivars with the conventional cultivating methods having in mind the teaching of D1.

In view of the fact that chicory roots are commonly used to produce inuline and that purified inulin, a product which should possibly not contain much side products such as monomeric saccharides, it is obvious for the man skilled in the art to choose chicory roots with higher DP and apply the known purification techniques (such as for example disclosed in EP-A-787 745, the claims). Claims 9 to 28 relating to conventional manufacture processes for extracting inulin and to the obtainable product do thus not meet the requirements of article 33(3) PCT.

The claimed subject matter is evidently industrially applicable.



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP99/00155

ad VIII:

Claims 25 and 26 do not fulfil the requirements of article 6 PCT since on the claimed products it is not verifiable whether they have been obtained by any known production process or by the processes claimed in claims 17 to 20.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C08B 37/18, A01G 1/00	A1	(11) International Publication Number: WO 99/37686 (43) International Publication Date: 29 July 1999 (29.07.99)
(21) International Application Number: PCT/EP99/00155 (22) International Filing Date: 13 January 1999 (13.01.99) (30) Priority Data: 98870012.6 21 January 1998 (21.01.98) EP (71) Applicant (for all designated States except US): TIENSE SUIKERRAFFINADERIJ N.V. [BE/BE]; Tervurenlaan 182, B-1150 Brussel (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): SMITS, Georges [BE/BE]; Dr. De Cockstraat 16, B-9308 Gijzegem (Aalst) (BE). DE LEENHEER, Leen [BE/BE]; Jezus Eiklaan 64, B-3080 Tervuren (BE). (74) Agent: HERMANS, Johnny; Tiense Suikerraffinaderij N.V., Aandorenstraat 1, B-3300 Tienen (BE).		(81) Designated States: AU, CN, IN, TR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS FOR THE MANUFACTURE OF CHICORY INULIN, HYDROLYSATES AND DERIVATIVES OF INULIN, AND IMPROVED CHICORY INULIN PRODUCTS, HYDROLYSATES AND DERIVATIVES (57) Abstract <p>The invention provides an improved process for the manufacture of chicory inulin, hydrolysates and derivatives of inulin by conventional manufacturing techniques from roots of chicory grown in appropriate regions and which have been grown and processed under proper climatological temperature conditions. Selection of said proper conditions enables to provide a growing and/or processing period for the chicory roots which may partly or wholly extend beyond the conventional periods. The invention relates also to improved grades of chicory inulin, of hydrolysates of chicory inulin, such as e.g. a polydisperse oligofructose composition and a fructose composition, as well as to the use of these products in the manufacture of food, feed, drinks, prophylactic and therapeutical compositions, chemical derivatives and non-food compositions. Improved standard grade chicory inulin according to the invention presents a degree of polymerisation (DP) which is at least 20 % higher than the one of conventional standard grade chicory inulin.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR THE MANUFACTURE OF CHICORY INULIN,
HYDROLYSATES AND DERIVATIVES OF INULIN, AND IMPROVED
CHICORY INULIN PRODUCTS, HYDROLYSATES AND DERIVATIVES.

5 Field of the invention

The present invention relates to a process for the manufacture of chicory inulin, hydrolysates and derivatives of inulin, to the use of certain chicory roots in such process, as well as to improved chicory inulin products, hydrolysates and derivatives of chicory inulin.

10

Background and prior art

Inulin is a carbohydrate which occurs in many plants and which can be produced by certain bacteria.

15

Inulin from plant origin consists of a polydisperse composition of chains of oligo- and polysaccharides which are composed of fructose units linked to each other through $\beta(2-1)$ fructosyl-fructose linkages, and which mostly terminate in one glucose unit. Inulin from plant origin is usually composed of linear chains, but may also contain some branched chains.

20

Main plant sources for inulin are roots of Chicory (*Cichorium intybus*) and tubers from Dahlia and Jerusalem artichoke, in which inulin can be present, respectively, in concentrations of about 15 - 18%, 12% and 14 to 18 % on fresh weight. Inulin can be readily extracted from these plant parts, purified and optionally fractionated to remove impurities, mono- and disaccharides and undesired oligosaccharides, in order to provide various

25

Inulin can be generally represented, depending from the terminal carbohydrate unit, by the general formulae GF_n and F_m , wherein G represents a glucose unit, F represents a fructose unit, n is an integer representing the number of fructose units linked to the terminal glucose unit, and m is an integer representing the number of fructose units linked to each other in the carbohydrate chain.

30

35

The number of saccharide units (fructose and glucose units) in one molecule, i.e. the values $n+1$ and m in the above formulae, are commonly referred to as the degree of polymerisation, represented by (DP). Often also the parameter (number) average degree of polymerisation,

represented by (\overline{DP}) , is used, which is the value \overline{DP}_n calculated, after complete hydrolysis, as follows:
$$\overline{DP}_n = \frac{\text{total}\%F}{\text{total}\%G} + 1$$

as in native inulins the F_m fraction is to be considered as negligible (in the equation % refers to % w/w). Furthermore, in this calculation the saccharides glucose (G), fructose (F) and saccharose (GF) which are present in the polydisperse polysaccharide, should not be taken into account, and the average degree of polymerisation is thus the (\overline{DP}_n) of inulin, herein interchangeably referred to in short as (\overline{DP}) inulin or (\overline{DP}) [cf. L. De Leenheer in "Production and use of inulin"; Carbohydrates as Organic Raw Materials, Vol. III, p. 67-92, (1996)].

The polysaccharide chains of native inulin from plant origin generally have a degree of polymerisation (DP) ranging from 3 to about 100, whereas the (\overline{DP}) largely depends from the plant source, the growth phase of the plant, the harvesting time, the storage conditions, and the process conditions used for the extraction and purification of the native inulin.

By the term native inulin is meant herein inulin that has been extracted from fresh plants or parts of plants, without applying any process to increase or decrease the (\overline{DP}) , while taking precautions to inhibit the plant's own hydrolase activity (e.g. by briefly boiling the extract) and to avoid hydrolysis. The (\overline{DP}) of native inulin thus essentially corresponds to the (\overline{DP}) of the inulin as present in the plant or plant parts.

The inulin obtained from plants or plant parts through conventional manufacturing techniques, including extraction, purification and isolation, without any process to increase the (\overline{DP}) of the native inulin, is termed herein, interchangeably, standard (\overline{DP}) grade inulin or standard grade inulin. As a consequence of the manufacturing process, the (\overline{DP}) of standard grade inulin is usually about 1 to 1,5 lower than the (\overline{DP}) of the corresponding native inulin.

With respect to the average degree of polymerisation (\overline{DP}) of standard grade chicory inulin, a distinction has to be made between the punctual (\overline{DP}) , which is the (\overline{DP}) of the inulin obtained from chicory roots processed at a specific moment of the growing or processing period, and the mean (\overline{DP}) , which is the mean value of the (\overline{DP}) of the inulin obtained from chicory roots, taken over a predetermined processing period adjacent to the growing period. By the processing period of the roots is meant herein the

period adjacent to the growing period up to the end of the processing of the chicory roots into inulin in a particulate form or up to a step in the inulin production where the intermediate obtained can be stored without degradation. Herein, the processing period thus includes the harvesting period which is the period, following the growing period, wherein no further inulin chain elongation (i.e. no further increase of (DP)) occurs but during which the roots still remain in the soil, as well as the harvesting as such, the possible storage period of the roots, which is preferably kept small, as well as their processing as such into inulin or up to a step in the inulin production where the intermediate obtained can be stored without degradation, e.g. by spray drying. Accordingly, by the processing of the roots is meant herein the harvesting (including the harvesting period and the harvesting as such), the possible storage and processing as such, and, by roots which have been processed is meant herein the harvesting (harvesting period and the harvesting as such), the possible storage, and the processing as such, of the roots into inulin or into said intermediate.

As a result of the standardisation sought for commercial products, the value of the (\overline{DP}) of commercial standard grade inulin commonly corresponds to the value of the mean (\overline{DP}) taken over the whole of the processing period, usually a period of about 60 to about 90 days.

The degree of polymerisation (DP) of native chicory inulin ranges from 3 to about 70. The punctual (\overline{DP}) of native inulin from chicory roots, which have been grown and processed under known, conventional conditions, ranges from about 10 to 14 (L. De Leenheer, o.c. , p. 71), whereas the mean (\overline{DP}) of known standard grade chicory inulin, i.e. inulin obtained from chicory roots which have been grown and processed under known, conventional conditions, is ranging from about 9 to 11, and is typically about 10.

Inulin of general formulae GF_n and F_m having a lower degree of polymerisation, usually defined as a $(DP) < 10$, is commonly named oligofructose and is referred to hereinafter accordingly.

Inulin is commercially available. For example, inulin from chicory is available under the brand name RAFTILINE® from ORAFTI®, (Belgium). Typical RAFTILINE® products are, for example , ST, ST-Gel and GR (i.e. all standard (\overline{DP}) grade inulin but in a different physical form, with a mean (\overline{DP}) inulin of about 10 and containing as by-products in total maximum

about 8 % by weight glucose, fructose and sucrose); LS (i.e. a low sugar standard (\overline{DP}) grade inulin with a mean (\overline{DP}) inulin of about 10 but which contains as by-products in total less than 1 % by weight glucose, fructose and sucrose); and HP and HP-Gel (i.e. both high performance grade inulin but in a different physical form, consisting both of fractionated, high (\overline{DP}) grade inulin with a (\overline{DP}) inulin (not a mean (\overline{DP}) inulin) of at least 23 which are essentially free of the by-products glucose, fructose and sucrose).

Although inulin has become available at industrial scale only from about 1990 onwards, it has already found major industrial applications in the food industry and is increasingly used also in the non-food industry.

Inulin is readily dispersible in aqueous liquids, is poorly soluble in cold water but fairly soluble in hot water and shows a good thermal stability. Depending from the presence and concentration of by-products, in particular monosaccharides, disaccharides and oligosaccharides, inulin has a taste ranging from slightly sweet to practically taste free. Inulin in food preparations, may, for example, improve texture and mouthfeel, and contribute to give more body and a smooth, creamy texture in low-fat products. Inulin also improves the stability of emulsions, dispersions, mousses, foams and creams. Furthermore, as disclosed in EP-A- 0 607 187, inulin can form with water a stable, fat-like, creamy structure (when subjected to mixing or shearing forces within a certain range of concentration and temperature). Moreover, inulin itself, i.e. when not taking into account mono- and disaccharides which are possibly present as by-products in the polydisperse polysaccharide, has a low caloric value, because when ingested, it is not digested in the small intestine of humans and many vertebrate animals, but is almost quantitatively entering the large intestine where it is fermented by bacteria, in particular by bifidobacteria. In view of its unique properties, inulin is widely used as a low calorie sugar replacement, fat replacement and/or texturising agent in various food products, e.g. in chocolate and confectionery products, in baked goods and breakfast cereals, in dairy products, such as desserts, creams, milks, cheeses, and yoghurts, in table spreads, in salad dressings, in fruit preparations, in frozen desserts, in meat products such as e.g. pâtés, and in drinks, as for example disclosed in EP-A- 0 599 830, EP-A- 0 607 187, EP-A- 0 607 189, and EP-A- 0 651 614. Moreover, inulin is also widely used in various feed products.

Furthermore, inulin which is a natural dietary fibre, has shown to have desirable health effects on humans and many vertebrate animals, such as e.g. beneficial prebiotic effects, in view of which inulin is already widely used in foods and drinks to improve the functioning of the digestive system, including stimulation of the grow, metabolism and activity of bifidobacteria while reducing those of less desirable intestinal bacteria.

Inulin has also been disclosed to have beneficial effects on lipid metabolism and to have prophylactic and therapeutic effects, e.g. on the genesis and development of breast cancer (EP-A-0 692 252) and of colon cancer in humans and non-bovine mammals (EP 97870069.8).

Furthermore, large quantities of inulin are used as raw material for the manufacture of oligofructose and fructose, products which are widely used in the food industry, e.g. in the manufacture of dairy products and drinks.

Besides, inulin is used as raw material for the manufacture of various chemical derivatives.

At present it is generally accepted that, compared to the actual applications and amounts used, inulin as such and as a raw material has still a much larger potential.

The strength of most technological and nutritional properties and effects of inulin, as well as its suitability as a raw material for the manufacture of oligofructose, fructose and inulin derivatives, are mostly proportional to its average degree of polymerisation (\overline{DP}): the higher the (\overline{DP}), the better.

Various processes for the manufacture of inulin from plant sources, including extraction, purification and fractionation, have already been disclosed.

In US 4.285.735, W.A. Mitchell et al. disclose a process for the manufacture of crystalline inulin from dahlia tubers which comprises a first isolation step wherein the tubers are finely ground and the water insolubles are removed from the water solubles; a next step wherein the obtained aqueous phase is boiled for a few minutes, followed by removal of the coagulated matter, concentration of the clarified solution to 40 - 70 % dry substance content, seeding with inulin crystals, cooling and storing for about 12 hours at 1° to 15° C; and a final step, in which the resulting crystalline inulin is isolated from the mother liquor and dried. No yield or average degree of polymerisation of the dry crystalline inulin has been disclosed.

In WO 94/12541 to Raffinerie Tirlemontoise S.A., a process is disclosed for removing to a large extent glucose, fructose and sucrose from standard grade chicory inulin via column chromatography techniques. The process involves charging a column containing a cation exchange resin with an aqueous, meta-stable, inulin solution containing 20 - 40 weight % dry substance, followed by elution with water preferably at 65 - 75°C, and yields inulin with a total concentration of said sugars below 1 weight %. This low sugar standard grade inulin has a (\overline{DP}) of about 10, which essentially corresponds to the (\overline{DP}) of the source chicory inulin.

Since many properties of inulin are favourably influenced by an increase of its average degree of polymerisation, various processes have already been developed for the preparation of inulin from plant sources with a (\overline{DP}) which is higher than the (\overline{DP}) of the corresponding native inulin.

One method, mentioned in International patent application WO 96/01849, consists in the fractionated precipitation of inulin from an aqueous solution by gradual decreasing the solubility of inulin in the solution through the addition of a lower alcohol. The longer polysaccharide chains precipitate before the shorter ones and can be separated from the latter by isolating them from the mixture, optionally in several fractions, by filtration or centrifugation. Accordingly, a rather pure product with a higher (\overline{DP}) than the (\overline{DP}) of the source inulin is obtained. However, the process is complex and laborious and the yield is very low, which makes the method technically and economically not attractive for a manufacture at industrial scale.

Furthermore, inulin with a higher (\overline{DP}) than native or source inulin can be prepared by classic fractional crystallisation techniques. Such method is for example described in Le Sillon Belge, April 24, (1989), according to which a warm aqueous polydisperse inulin solution is gradually cooled using ceiling temperatures between 40°C and 10°C, where necessary under grafting with inulin crystals. Since inulin molecules with a higher degree of polymerisation are less soluble than the ones with a lower degree of polymerisation, the former will precipitate before the latter, thus enabling to separate them from each other and to isolate them from the mixture by filtration or centrifugation. The method being laborious, is economically not attractive and hardly suitable for manufacture at industrial scale.

A similar method has been described by E. Berghofer (Inulin and inulin containing crops, Ed. A. Fuchs, Elseviers Sc. Publ., p. 77, (1993), according to which an aqueous solution of polydisperse chicory inulin is subjected to crystallisation with a slow cooling pattern (3°C/hour from 95°C to 4°C) under addition of grafting crystals. The longer inulin chains precipitating before the shorter ones, the former can be separated from the latter and are isolated by filtration or centrifugation. The method, however, suffers from disadvantages which are comparable to the ones of the method disclosed in Le Sillon Belge, indicated above.

According to a further method, described in EP 0 627 490 to Südzucker A.G., polydisperse native inulin is treated with inulinase enzyme to break down the inulin fraction with a low degree of polymerisation, thus increasing the (\overline{DP}) of the remaining inulin fraction. The degradation products formed, essentially glucose and fructose, have to be removed by a supplementary purification step, which forms a technical and economical drawback.

In WO 96/01849 to Tiense Suikerraffinaderij, an economically more attractive process is described for the manufacture at industrial scale of inulin with an average degree of polymerisation which is almost double or even higher, than the (\overline{DP}) of the source inulin. The process involves a directed crystallisation of inulin from a meta-stable aqueous solution of a polydisperse source inulin, yielding a suspension of crystallised inulin particles. Typically said source inulin is crude chicory inulin, i.e. inulin extracted from roots which has subsequently been subjected to a depuration treatment (i.e. a treatment involving liming, carbonatation and filtration) or standard grade chicory inulin with a (\overline{DP}) of about 10. The inulin particles formed are isolated by filtration or centrifugation. Before drying, they are washed with cold, demineralised water which removes most of the remaining impurities, mono- and disaccharides, as well as most of the remaining lower oligomeric polysaccharides ($DP < 10$) from the particles. In this way, from a source chicory inulin with a (DP) ranging from 3 to about 70 and a (\overline{DP}) of about 10, high performance chicory inulin is obtained having a (DP) ranging from about 10 to about 70 and a (\overline{DP}) of about 20 to about 30 with an average yield of about 30 to about 35%. The isolated crystalline inulin can optionally be transformed into a powder form through spray drying techniques.

In view of its natural origin, its desirable technological properties and the fact that it is a dietary fibre with beneficial health effects on humans and many vertebrate animals, there is a steadily increasing demand in the food, the feed and the pharmaceutical industries, as well as in the non-food industries, for inulin, in particular for inulin grades with a higher (\overline{DP}).

Accordingly, industry is steadily facing the problem of providing technically and economically attractive methods and processes enabling the manufacture of inulin at industrial scale, preferably with improved yield and/or improved characteristics and properties, as, for example, an increased average degree of polymerisation.

The extraction of inulin from chicory roots, followed by purification, conventionally including depuration and refining (i.e. involving treatment over ion-exchangers, treatment with active carbon and filtration), optionally followed by fractionation, and isolation in particulate form of the obtained inulin, constitutes the most important manufacturing method of inulin at industrial scale. This is a result of the fairly availability of chicory roots, the inulin content of the roots, and the suitability of the roots for processing at large scale. The manufacturing techniques commonly used are well known in the art.

Chicory is conventionally cultivated in certain northern parts of Western Europe, where it is seeded in Spring (usually in April) and the roots are harvested, stored and processed for inulin production late Autumn, usually from about mid September to about end November, yielding through conventional manufacturing techniques standard grade chicory inulin with a mean (\overline{DP}) of about 10. The whole growing and processing period covers about 150 to about 230 days. It is known that the degree of polymerisation (DP) and the punctual (\overline{DP}) of the inulin, as well as the content of inulin in the chicory roots (i.e. the % by weight of inulin in the fresh root material) are increasing during the growing season to reach a maximum after about 150 days of growing, whereas the biomass of the roots and thus the yield (in ton/ha) of inulin increase till about the end of the growing season. The end of the growing season is the time when the biomass of the roots ceases to increase significantly (i.e. after about 180 to about 200 days of growing), which usually corresponds to the end of October. Towards the end of the growing season, the (DP) and punctual (\overline{DP}) of the inulin in roots remaining in the soil, as well as in harvested and stored

roots, are decreasing with time, and the rate of decrease is usually significantly increased from about begin November. This situation considerably limits the periods for growing (including seeding and growing) and for processing, including harvesting (harvesting period and harvesting as such), storage, and processing as such, of the chicory roots for the production of inulin, hydrolysates and derivatives of inulin in a technically and economically attractive manner.

Consequently, in spite of the fact that the manufacture of inulin from chicory roots constitutes the most important route to inulin, said manufacture is nevertheless confronted with considerable hurdles and problems, including: (i) the limited period, including limitations in duration of the period as well as in time period of the year, during which chicory roots can be seeded, grown, harvested, stored and processed for the manufacture of inulin in a technically and economically attractive manner and/or without undergoing a significant decrease of the (\overline{DP}) of the inulin in the roots, (ii) the rather low mean (\overline{DP}) of standard grade chicory inulin (having a (\overline{DP}) of about 10), (iii) the need to include a fractionation step in the manufacturing process of chicory inulin when, e.g. for technological or nutritional reasons, inulin is required with a (\overline{DP}) which is higher than the (\overline{DP}) of about 10 of known standard chicory inulin, (iv) the rather poor yields of the known fractionation processes leading to inulin with a higher (\overline{DP}) when chicory inulin of standard grade (with a mean (\overline{DP}) of about 10) is used as the source inulin, and (v) the economical unattractive situation of the plants for the processing of the chicory roots. These plants are commonly operated during only a few months a year and are sized to process a large quantity of chicory roots in a very limited period. The processing as such of the roots for the manufacture of inulin, or of an intermediate as mentioned above, commonly takes about a day, whereas the processing period, including the harvesting (harvesting period and harvesting as such), storage and processing as such, of the roots, commonly lasts about 60 to about 90 days.

Objects of the invention

The present invention aims to solve one or more of said problems, inter alia, by providing a new process for the manufacture of chicory inulin. In particular, the invention aims to provide a process for the manufacture of chicory inulin, partial hydrolysates, complete hydrolysates and

derivatives of inulin, from chicory roots with a growing and/or a processing period which partially or wholly falls outside the conventional ones. Furthermore, the invention aims to provide a process for the manufacture of standard grade chicory inulin with an increased mean (\overline{DP}) compared to known standard grade chicory inulin while possibly providing additional technical and economical advantages, to provide an improved standard grade chicory inulin and uses thereof, as well as to provide a process for the manufacture of high performance grade chicory inulin with increased yield. Furthermore, the invention aims to provide an improved process for the manufacture of hydrolysates and derivatives of chicory inulin.




Description of the invention

The production of inulin in chicory is an enzymatically driven process wherein the so-called SST gene and FFT gene of the chicory genome are involved.

The SST gene codes for a transferase enzyme that catalyses the transfer reaction of fructose from sucrose synthesised by the chicory plant, to form a lower-(DP) oligofructose (mainly GF₂), as schematically represented in equation (1) below, wherein G, F and GF respectively represent a glucose, fructose and sucrose unit, and GFF represents the lower-(DP) oligofructose GF₂:





The FFT gene codes for a transferase enzyme that catalyses the transfer of fructose from one oligofructose, usually a GF₂ oligofructose, to another, which results in the increase by one unit of the (DP) of the oligofructose substrate, finally resulting in inulin chains of the general formula GF_n with a (DP) > 10, as schematically represented in equation (2a, 2b, and 2c) below, wherein G, F, n and GFF have the meaning defined above and GFFF, GFFFF, GF_{n-1} and GF_n represent an oligosaccharide or polysaccharide (inulin) depending on the value of n :



On the other hand, it is known that inulin chains can also undergo degradation in chicory roots subsequently to their synthesis.

It has now been found that the degradation is mainly catalysed by one or more enzymes with exohydrolase activity (hereinafter referred to as exohydrolase enzyme) coded for by one or more FEH genes (hereinafter referred to as FEH gene) of the chicory genome.

5 The degradation which occurs through a repetitive reaction, can be schematically represented by equation (3a) and (3b) below, wherein G, F, GF_n and n have the meanings defined above:



10 The inventors have now found that the activity of the FEH gene in the chicory plant, and thus the production of exohydrolase enzyme and the corresponding degradation of inulin chains in the chicory roots, are essentially triggered by the occurrence of conditions of low temperature during the processing period and during a large part of the growing period of the roots, particularly from the beginning of the third month of the growing period onwards.

15 By said conditions of low temperature are meant not only those temperature conditions at which the chicory roots become frozen (i.e. conditions at which plant cells in the roots are destroyed), but also those at which the roots become frost-bitten (i.e. conditions at which the cells in the roots are not destroyed, but at which, as a consequence of the activity of the FEH gene, an altered metabolism starts off according to which the roots start to metabolise the inulin molecules). Said low temperature conditions are, in particular, conditions corresponding to a temperature in thermometer shelter below about minus 1°C which corresponds to a temperature of about minus 3°C to minus 5°C at the soil surface.

25 It can be found out easily whether or not the FEH gene in the chicory roots has been triggered because triggering of the FEH gene is marked by a significant increase in concentration of fructose in the chicory roots (see equation (3a) and (3b) above).

30 Taking the above into account, the inventors have found that in appropriate regions for growing chicory which in addition have the proper climatological temperature conditions, chicory roots can be grown and processed into inulin or into hydrolysates or derivatives of inulin, outside the known, conventional growing and processing periods. They also found

35

that improved chicory inulin products, particularly improved standard grade chicory inulin, can be obtained as well as improved hydrolysates and derivatives of said inulin, from chicory roots which have been grown and processed in said regions under said proper climatological temperature conditions.

By appropriate regions is meant herein regions which present the known conditions for growing chicory, such as, for example, conditions regarding soil and humidity.

By proper climatological temperature conditions are meant conditions which are such that the FEH gene in chicory roots is not triggered by the occurrence of low temperature conditions as defined above.

Based on the above findings, but not wishing to be bound by any theory set forth above, there is provided by the present invention a solution to one or more of the above mentioned problems regarding the provision of chicory inulin, hydrolysates and derivatives of inulin.

Accordingly, in a first embodiment of the invention, a process is provided for the manufacture of chicory inulin, hydrolysates of inulin or derivatives of inulin, from chicory roots through conventional manufacturing techniques, wherein the source material for the process are roots of chicory grown in appropriate regions and which have been grown and processed under proper climatological temperature conditions which are such that during a period of at least from the beginning of the third month of the growing period till the end of the processing of the chicory roots the FEH gene in the chicory roots has not been triggered (i.e. not triggered at all or not triggered to a significant extent) by the occurrence of low temperature conditions, said chicory roots have had a growing period of at least 150 days, preferably of at least 160 days, more preferably of about 180 days, and said chicory has been seeded in the northern hemisphere within a period selected from the periods ranging from December 1 till March 14, from May 15 till May 31, and from June 1, preferably from June 15, till November 30, provided that when said chicory has been seeded in the period from May 15 till May 31, or from June 1 till June 14, the chicory roots have had a growing period of at least 180 days, or in the southern hemisphere within a period selected from the periods ranging from June 1 till September 14, from September 15 till September 30, from October 1 till

November 14, from November 15 till November 30, and from December 1, preferably from December 15 till May 31.

5 In a preferred embodiment, the climatological temperature conditions are such that the FEH gen is not triggered neither during a processing period of 60 days, more preferably 90 days, adjacent to the growing period.

In a preferred embodiment, the chicory inulin is standard grade chicory inulin with a (\overline{DP}) of at least about 10, low sugar standard grade chicory inulin with a (\overline{DP}) of at least about 10, or high performance grade chicory inulin with a (\overline{DP}) of at least about 20

10 In a further preferred embodiment, the chicory inulin is improved standard grade chicory inulin (which is standard grade chicory inulin with a mean (\overline{DP}) of at least 12, more preferably of at least 14), improved low sugar standard grade chicory inulin (which is low sugar standard grade chicory inulin with a mean (\overline{DP}) of at least 12, preferably of at least 14), or improved
15 high performance grade chicory inulin (which is high performance grade chicory inulin with a mean (\overline{DP}) of at least 20, preferably at least 23, more preferably at least 25), the mean (\overline{DP}) being taken over a processing period of at least 60 days. To prepare inulin with a mean (\overline{DP}) being taken over a certain processing period, the batches of inulin obtained during said period
20 are combined, for example by blending.

In a further preferred embodiment, the process yields a partial hydrolysate of inulin being preferably an oligofructose composition or a complete hydrolysate of inulin being a fructose composition, in particular a fructose rich fructose syrup.

25 The inventors have furthermore found that, in appropriate regions, under proper climatological temperature conditions, chicory roots can be grown and processed into inulin, including a growing period of the chicory of at least 160 days, preferably of about 180 days, and a processing period of the roots of at least 60 days, preferably of about 90 days, without that
30 considerable inulin chain degradation occurs in the roots, and that from chicory roots grown and processed under these proper climatological temperature conditions, improved standard grade chicory inulin, can be obtained through conventional manufacturing techniques, without any fractionation, with a mean (\overline{DP}) taken over a processing period of at least 60
35 days, preferably of 90 days, which is at least 20% higher, usually from 30 to 50

% higher, and typically about 40% higher, than the mean (\overline{DP}) taken over a corresponding processing period of known standard grade chicory inulin.

By said proper climatological temperature conditions in this embodiment are meant conditions wherein the FEH gene in the chicory roots is not triggered (i.e. not triggered at all or not triggered to a significant extent) by the occurrence of said low temperature conditions within a period of at least 220, preferably at least 240, consecutive days immediately preceding the end of the processing of the roots.

By known standard grade chicory inulin is meant herein inulin obtained by conventional manufacturing techniques (without a process step to increase the (\overline{DP}) of the native inulin) from chicory roots which have been grown and processed under conventional climatological temperature conditions.

By conventional climatological temperature conditions is meant herein conditions wherein within a period of less than 220 consecutive days immediately preceding the end of the processing of the roots, low temperature conditions have occurred which triggered the FEH gene in chicory roots to a significant extent.

Whereas known standard grade inulin commonly has a mean (\overline{DP}) (over a processing period of at least 60 days) of about 10, improved standard grade inulin from chicory roots grown and processed under said proper climatological conditions has a mean (\overline{DP}) (over a processing period of at least 60 days), of at least 12, usually from 13 to 16, and typically of at least 14.

Accordingly, in a further embodiment of the present invention, a process is provided for the manufacture of improved standard grade chicory inulin from chicory roots through conventional manufacturing techniques, wherein the source material for the process are roots of chicory grown in appropriate regions and which have been grown and processed under proper climatological temperature conditions wherein, within a period of at least 220, preferably of at least 240, consecutive days immediately preceding the end of the processing of the roots, no low temperature conditions occurred which triggered the FEH gene in chicory roots, said roots have had a growing period of at least 160 days, the FEH gene in the chicory roots has not been triggered by the occurrence of low temperature conditions within the period from at least the beginning of the third month of the growing period till the end of the processing of the roots, and the product obtained is

improved standard grade chicory inulin with a mean (\overline{DP}) taken over a processing period of at least 60 days adjacent to the growing period, which is at least 12, usually from 13 to 16, and preferably at least 14.

5 In a preferred embodiment of said process, the climatological temperature conditions are such that no low temperature conditions which trigger the FEH gene in chicory roots occur within a total period of at least 270 consecutive days, more preferably of at least 300 consecutive days, immediately preceding the end of the processing of the roots, and most preferably within a period covering a complete year.

10 In a further preferred embodiment of said process for the manufacture of improved standard grade chicory inulin, the chicory has been seeded in the northern hemisphere at least within a period selected from the periods ranging from December 1 till March 14, from March 15 till May 14, from May 15 till May 31, and from June 1 till November 30, or in the southern
15 hemisphere at least within a period selected from the periods ranging from June 1 till September 14, from September 15 till November 14, from November 15 till November 30 and from December 1 till May 31.

In the processes according to the invention the said proper climatological temperature conditions are preferably such that during the
20 concerned period immediately preceding the end of the processing of the chicory roots the temperature in the temperature shelter has not dropped below minus 1°C, which usually corresponds to a soil surface temperature of about minus 3° to minus 5°C. More preferably, the temperature in the temperature shelter has not dropped below 0°C, usually corresponding to a
25 soil surface temperature of about minus 2°C to minus 4°C. Most preferably, the temperature in the temperature shelter has not dropped below plus 3°C.

The inventors have found that appropriate regions for growing chicory which present, besides the known conditions for growing chicory, said proper climatological temperature conditions, do exist in certain regions of
30 the USA, Europe, Africa, South America, Australia and Asia. Typically, such appropriate regions with the proper climatological temperature conditions are found in the Californian region of the USA, in certain valleys of Chile, and in certain maritime regions of Europe, Turkey, China, India, Morocco, Algeria and South Africa.

35 Under the proper climatological temperature conditions in accordance with the present invention, the yield of chicory roots per hectare (ranging

from about 25 to 100 ton/hectare) as well as the total content of carbohydrates and the content of inulin in the roots (ranging, respectively, from about 14 % to about 20 % and from about 13 to about 19 % by weight of the fresh roots), are comparable to the ones observed for roots of chicory grown under conventional climatological temperature conditions.

The above process according to the invention does not only provide a new, improved standard grade chicory inulin, having improved properties resulting from its higher (\overline{DP}) compared to known standard grade chicory inulin, but also provides improved intermediates or source material, such as depurated inulin, refined inulin or purified inulin solutions or concentrates, for the manufacture according to conventional techniques of special grades of inulin, such as improved low sugar standard grade chicory inulin and improved high performance grade chicory inulin, hydrolysates and derivatives of inulin.

The various process steps of the above processes for the manufacture of standard grade, improved standard grade, low sugar standard grade, improved low sugar standard grade, high performance grade, and improved high performance grade chicory inulin, of partial and complete hydrolysates and derivatives of standard grade and of improved grade chicory inulin, can be carried out by techniques which are well known in the art, some of which correspond well to techniques commonly used in the manufacture of sugar from sugar beets or cane.

In a typical execution form of said processes for the manufacture of, respectively, standard grade chicory inulin and improved standard grade chicory inulin, the conventional manufacturing techniques comprise the following consecutive steps:

(i) isolation of the inulin from the chicory roots, providing an aqueous solution of crude inulin;

(ii) purification of the crude inulin obtained in step (i), yielding an aqueous solution of purified inulin, optionally followed by concentration of this solution by partial removal of the water yielding purified inulin concentrate; and

(iii) isolation of the inulin from the aqueous solution or concentrate of purified inulin obtained in step (ii), yielding respectively standard grade chicory inulin and improved standard grade chicory inulin, in particulate form.

Suitable techniques for the isolation of inulin from chicory roots comprise, for example, grinding and wet milling or crushing of the chicory roots, or, preferably, extraction of the inulin from freshly sliced or shredded chicory roots with hot water.

5 Suitable techniques for the purification of the crude inulin obtained in step (i) comprise, for example, depuration of the aqueous liquid obtained in step (i) (involving e.g. liming and carbonatation or another flocculation technique, and filtration), followed by refining (involving e.g. treatment over ion-exchangers, treatment with active carbon and filtration, or
10 ultrafiltration), whereas the solution can be concentrated by partial removal of the water by means of heat and/or reduced pressure. The isolation of the standard grade chicory inulin or improved standard grade chicory inulin in particulate form from the aqueous solution or concentrate of purified inulin obtained in step (ii) above is typically carried out by spray drying, as for
15 example described in WO 94/19973.

In a further aspect of the invention, a process is provided for the manufacture of improved low sugar standard grade chicory inulin, containing in total less than 1 weight % monomeric saccharides and sucrose, wherein improved standard grade chicory inulin obtained in step (iii) of the
20 above process or a solution or concentrate of purified inulin obtained in step (ii) of the above process, is used as a source material, and is, in accordance with known manufacturing techniques, subjected to the following consecutive steps:

(iv) removal of the monomeric saccharides and sucrose, yielding
25 a low sugar standard grade inulin solution or concentrate, and
(v) isolating the low sugar standard grade inulin in particulate form from the solution or concentrate obtained in step (iv),
yielding improved low sugar standard grade chicory inulin with a mean (\overline{DP}), taken over a processing period of at least 60 days, of at least 12,
30 typically of 13 to 16, preferably of at least 14.

Said removal of monomeric saccharides and sucrose can be carried out, for example as described in PCT patent application WO 94/12541, through column chromatography of an aqueous, meta-stable solution of the source inulin, whereas the improved low sugar standard grade chicory inulin
35 obtained can be isolated in particulate form through spray drying, for example as described in PCT patent application WO 94/19973.

Moreover, the inventors also have found that starting from improved standard grade chicory inulin or from its intermediate, depurated inulin or purified inulin solution or concentrate obtained in step (ii) of the above process according to the invention, and having a (\overline{DP}) of at least 12, the yield of high performance grade chicory inulin obtained through conventional fractionation techniques is significantly increased compared to the yield obtained by said conventional techniques when known standard grade inulin or its intermediate, depurated or refined inulin solution or concentrate, with a (\overline{DP}) of about 10, is used as a source material.

By high performance grade chicory inulin is meant herein inulin which has been subjected to one or more fractionation processes to increase the (\overline{DP}) of the source inulin in order to reach a (\overline{DP}) of at least 20.

Whereas commonly a yield of high performance grade inulin of about 30 % to about 35 % is obtained (calculated on the dry substance of the source material) through conventional techniques starting from known standard grade chicory inulin or the corresponding intermediate depurated or refined chicory inulin with a (\overline{DP}) of about 10, as e.g. described in PCT patent application WO 96/01849, a yield of 40 % or more of high performance grade inulin of a (\overline{DP}) of at least 20 is obtained by said conventional techniques when improved standard grade inulin or its intermediate, depurated or refined inulin obtained in the process according to the present invention with a (\overline{DP}) of at least 12, preferably of at least 14, is used as a source material.

Accordingly, in a further aspect of the subject invention, an improved process is provided for the manufacture of high performance grade chicory inulin with a (\overline{DP}) of at least 20 which is essentially free from low molecular monomeric saccharides, dimeric saccharides and oligofructose, and essentially free from colourings, salts, proteins, organic acids and technological aids such as solubility affecting products, wherein improved standard grade chicory inulin or its intermediate, depurated or refined inulin solution or concentrate, obtained in step (ii) of the above process according to the invention, with a (\overline{DP}) of at least 12, is used as a source material and is subjected in accordance with known manufacturing techniques to the following consecutive steps:

(vi) fractionation , and

(vii) isolation in particulate form of the high performance grade inulin from the fractionated product obtained in step (vi), thereby yielding high performance grade chicory inulin with a (\overline{DP}) of at least 20 in a yield of 40 % or more based on the source inulin.

5 In a preferred embodiment, the high performance grade inulin is obtained in a yield of 45 % and has a (\overline{DP}) between 20 and 35, preferably of at least 23.

10 In another preferred embodiment of the above process, the source material, improved standard grade chicory inulin or an intermediate thereof, such as depurated or refined chicory inulin, has a mean (\overline{DP}) of at least 14, and the product obtained is improved high performance grade chicory inulin, which is obtained in a yield of more than 40%, preferably at least 45%, typically between about 45% and 50%, based on the source material, with a mean (\overline{DP}), taken over a processing period of at least 60
15 days, of at least 23, preferably of at least 25.

The steps of the above process for the manufacture of high performance grade or improved high performance grade chicory inulin can be carried out according to conventional techniques which are well known in the art. The fractionation is preferably carried out through directed
20 crystallisation of an aqueous metastable inulin solution, yielding an aqueous suspension of fractionated inulin. The isolation of the fractionated inulin in particulate form from said suspension is commonly carried out by filtration, centrifuging or the like techniques, usually followed by washing with water and drying of the isolated particles, as e.g. described in PCT Patent
25 Application WO 96/01849. When depurated inulin is used as a source material, it may be beneficial to include a refining step in the fractionation process.

Optionally, the obtained high performance grade or improved high performance grade inulin can be transformed into a powder form composed
30 of particulates and/or aggregates or conglomerates, by thinning the isolated high performance grade inulin with water followed by spray drying according to known techniques, as for example described in PCT patent applications WO 94/19973 and WO 96/01849.

35 Furthermore, chicory inulin obtained according to the present invention can optionally be associated in the isolation step of the inulin

particulates, or in an additional step, to other food or feed components, such as, for example, sucrose, high intensity sweeteners, dairy, and extracts of coffee, cacao or tea, through known techniques, for example by co-spray drying as described in EP 96870099.7.

5 In a further aspect of the present invention, an improved process is provided for the manufacture of a polydisperse oligofructose composition, by known techniques from inulin, e.g. by partial, acidic or enzymatic, hydrolysis of inulin and subsequent purification, for example as described in
10 patent application PCT/BE 97/00087, wherein improved standard grade chicory inulin with a (\overline{DP}) of at least 12, preferably of at least 14, or its intermediate, depurated or refined inulin solution or concentrate, obtained in a process according to the present invention described above, is used as a source material, and the obtained polydisperse oligofructose composition has a (DP) ranging from 2 to 10 and contains at least 90%, calculated on the
15 dry substance, polysaccharide chains of formula F_m and GF_n wherein G and F have the meaning of a glucose unit and fructose unit, respectively, and n and m are integers.

In a further aspect of the subject invention, an improved process is provided for the manufacture according to known techniques of a fructose
20 composition by complete hydrolysis of chicory inulin, e.g. by acidic or enzymatic hydrolysis, followed by purification, for example, as described in patent application EP 97870111.8, wherein improved standard grade chicory inulin with a (\overline{DP}) of at least 12, preferably of at least 14, or its intermediate, depurated or refined inulin solution or concentrate, obtained in the above
25 process according to the present invention, is used as a source material, thereby yielding directly, i.e. without a fractionation step, a fructose composition, for example a fructose syrup, containing at least 89 % by weight of fructose, calculated on dry substance. In the obtained fructose composition the ratio F/G is higher than in a fructose composition obtained by
30 conventional techniques from known standard grade inulin or from its intermediate, depurated or refined inulin solution or concentrate.

In still another aspect of the subject invention, a process is provided for the manufacture according to known techniques of improved derivatives of inulin, for example by etherification, esterification, carboxymethylation,
35 ethoxylation with ethylene oxide, cationisation or cross-linking, wherein improved standard grade chicory inulin or its intermediate, depurated or

refined inulin solution or concentrate, or high performance grade inulin, obtained in the process according to the present invention described herein before, is used as a source material.

According to another embodiment, the present invention relates to improved standard grade chicory inulin with, expressed on dry matter, an inulin content of at least about 92% by weight and a maximal total content of glucose, fructose and sucrose of 8 % by weight, and with a mean (\overline{DP}) (after processing) - taken over a processing period of the chicory roots used as source material in the manufacture process of said inulin of at least 60 days, preferably of at least 90 days - of at least 12, typically of 13 to 16, preferably of at least 14.

In a further aspect, the invention relates to improved low sugar standard grade chicory inulin with, expressed on dry matter, an inulin content of more than 99% by weight and a total content of glucose, fructose and sucrose of less than 1% by weight, which has a (\overline{DP}) of at least 12, typically of 13 to 16, preferably of at least 14.

In still a further aspect, the invention relates to improved high performance grade chicory inulin which is essentially free from low molecular monomeric saccharides, dimeric saccharides and oligofructose, and essentially free from colourings, salts, proteins, organic acids and technological aids such as solubility affecting products, with a mean (\overline{DP}) , taken over a processing period of the source chicory roots of at least 60 days, of at least 20, preferably at least 23, and more preferably of at least 25.

The said improved grades of chicory inulin are preferably present in the form of spray dried particulates or of spray dried aggregated and/or agglomerated particles, and are obtainable by the above described manufacturing processes according to the present invention.

In still a further aspect, the invention relates to an improved polydisperse oligofructose composition, respectively to an improved fructose composition, obtainable by partial, respectively by complete hydrolysis of improved standard or high performance grade chicory inulin according to the present invention, as defined above.

In another embodiment, the present invention relates to the use of roots of chicory grown in appropriate regions as a source material in a process for the manufacture of chicory inulin, of partial hydrolysates of inulin, of complete hydrolysates of inulin, or of derivatives of inulin, said

roots having been grown and processed under proper climatological temperature conditions which are such that within a period of at least from the beginning of the third month of the growing period till the end of the processing period, the FEH gene in the chicory roots has not been triggered by the occurrence of low temperature conditions, said chicory roots have had a growing period of at least 150 days, preferably of at least 160 days, more preferably of at least 180 days, and said chicory has been seeded in the northern hemisphere within a period selected from the periods ranging from December 1 till March 14, from May 15 till May 30, and from June 1, preferably from June 15, till November 30, provided that when said chicory has been seeded in the period from May 15 till May 31, or from June 1 till June 14, the chicory roots have had a growing period of at least 180 days, or in the southern hemisphere within a period selected from the periods ranging from June 1 till September 14, from September 15 till September 30, from October 1 till November 14, from November 15 till November 30, and from December 1, preferably December 15, till May 31.

In a further embodiment, the present invention relates to the use of roots of chicory grown in appropriate regions as a source material in a process for the manufacture of chicory inulin, of partial hydrolysates of inulin, of complete hydrolysates of inulin, or of derivatives of inulin, said roots having been grown and processed under proper climatological temperature conditions which are such that within a period of at least 220 consecutive days, preferably at least 240 consecutive days, more preferably of at least 270 consecutive days, even more preferably of at least 300 consecutive days, and most preferably within a period covering a complete year, no low temperature conditions occur which trigger the FEH gene in chicory roots, and said roots have had a growing period of at least 160 days and preferably of at least 180 days, and no low temperature conditions which triggered the FEH gene in the roots occurred from the beginning of the third month of the growing period till the end of the processing period of the roots.

Typical processes for said manufacture and typical grades of chicory inulin, of partial and complete hydrolysates of inulin, are, for example, the ones described above.

In still another embodiment, the present invention relates to the use of improved standard grade chicory inulin, improved low sugar standard

grade chicory inulin and improved high performance grade chicory inulin according to the present invention in the manufacture of food, feed, and drinks, particularly in the manufacture of low calorie, dietary, functional, and prebiotic food compositions, feed compositions and drinks as well as in the manufacture of non-food compositions.

In still a further embodiment, the present invention relates to improved standard grade chicory inulin, improved low sugar standard grade chicory inulin and improved high performance grade chicory inulin according to the present invention for use in the manufacture of a prophylactic or therapeutic composition, for use as a medicament, and for use in a prophylactic and/or therapeutic method for treatment of humans and vertebrate animals, in particular for the stimulation of calcium-uptake, the increase of mineral bone density, for the prevention or treatment of osteoporosis and of cancer, typically breast cancer and colon cancer.

In an ultimate embodiment, the present invention relates to the use of improved standard grade chicory inulin, improved low sugar standard grade chicory inulin and improved high performance grade chicory inulin according to the present invention as a raw material in the manufacture of chemical derivatives of inulin, such as, for example, derivatives obtained by etherification, esterification, carboxymethylation, ethoxylation, cationisation and cross-linking.

As a source material for use in the various embodiments according to the present invention, preferably a variety of chicory is chosen which gives little bolting under the given growing conditions, most preferably a chicory variety which is conventionally classified as "bolting resistant", such as for example the variety "Orchies".

Furthermore, the chicory variety is preferably chosen conventionally while taking into account the particulars of the soil and the particulars of the climatological conditions of the area in which the roots have to be grown. In certain regions where humidity is low and/or the temperature is high compared to conventional regions where chicory is grown, the chicory roots are preferably cultivated using irrigation. The type of irrigation, such as for, example, furrow-, aspersion- or sprinkling-type irrigation, is preferably chosen in accordance with the kind of soil and/or the climatological conditions.

The processes for the manufacture of the various mentioned grades of chicory inulin, hydrolysates and derivatives of inulin, the improved products and their uses according to the present invention, as well as the use for the manufacture of said products of chicory roots which have been grown and processed under the proper climatological temperature conditions in accordance with the present invention as defined above, present various considerable technological, nutritional and/or economical advantages over the known processes and products.

For example, the invention enables the direct manufacture by conventional techniques of improved standard grade chicory inulin as well as of improved high performance grade chicory inulin, with a mean (\overline{DP}) taken over a predetermined processing period which is significantly higher than the mean (\overline{DP}) of the corresponding known grade of chicory inulin. This obviously results in various cost savings, because no fractionation step, respectively no additional fractionation step, is required to obtain said improved inulin, thus resulting in savings of time and manufacturing costs. Furthermore, often smaller quantities of the improved inulin products can be used compared to the corresponding known inulin products in given compositions to obtain a predetermined technological effect, such as, for example, a stabilising or texturising effect. This also results in product savings and in cost savings and provides an increased flexibility in the content of ingredients of, for example, food, feed and drinks compositions. Hydrolysates from improved standard grade inulin or its intermediate present a higher fructose content, respectively a higher oligofructose content, than the corresponding hydrolysates from known standard grade chicory inulin without an additional refining of the source product or the hydrolysates, which obviously results in advantages with respect to e.g. nutritional aspects, manufacturing time and costs.

Chemical derivatives from improved standard grade chicory inulin according to the present invention can show improved physico-chemical properties and/or improved functionality compared to corresponding derivatives obtained from known standard grade chicory inulin.

Besides, by the selection of an appropriate region with the proper climatological temperature conditions for growing and processing roots of chicory in accordance with the present invention, it is possible to extend the conventional growing and processing periods in duration as well as in time

period of the year. In regions with said proper climatological temperature conditions, chicory can not only be seeded or seedlings (transplants) planted in the beginning of Spring, but also at other points in time of the year, for example, even in Autumn. Accordingly, due to the non-triggering of the FEH gene under said proper climatological temperature conditions, the seeding period can be extended to a period of at least eight months compared to the conventional period of about 50 days. Also the harvesting, storage and processing of the chicory roots to produce inulin or hydrolysates is not limited any more to the end of Autumn, but can be spread over a large part of the year. The longer the period with the proper climatological temperature conditions, the longer the periods available for seeding, growing, harvesting, storage and processing of the chicory roots. In appropriately selected regions, it even becomes possible to seed, grow, harvest, store and process roots of chicory to produce inulin, hydrolysates and derivatives of inulin, around the year. This extended period for growing and processing chicory roots for the production of inulin, hydrolysates or derivatives of inulin, obviously results in significantly technical and economical advantages. For example, the supply of chicory roots for said production can be spread over large parts of the year, even over a complete year, and as a result thereof, the processing plants do not become confronted any more with one short period per year for the processing of the chicory roots. This makes it possible to down size the processing plants and/or exploit them more efficiently and in an economically more attractive manner.

25

The invention is illustrated by the examples given below.

Example 1 (Comparative):

Standard grade chicory inulin from chicory roots grown in Belgium under conventional climatological temperature conditions (fall harvest)

30

Chicory, variety Madonna, coated seeds, were seeded on 16/04 in Opheylissem (Belgium) and grown under non-irrigated conditions; population of 150.000 plants/ha.

35

Harvest was done manually two-weekly from 16/09 until 10/12 and the roots were processed without delay.

A. Extraction of inulin was done at laboratory scale from chicory gratings which were prepared by slicing the roots in pieces of about 2cm/1cm/0,5 cm and blending them in a Warring* (* trade name) blender up to complete homogenisation. The extraction was performed by treating the gratings with boiling distilled water (300 g water for 60 g gratings in a 500 ml Duran* (*trade name) bottle). To achieve complete extraction, the mixture of gratings and water is incubated in a closed bottle at 75 °C for 60 minutes with stirring at 135 rpm. The pH of the mixture is set between 5,5 and 7, if necessary, before incubation to avoid acid hydrolysis. Finally, a centrifugation of 10 min at 4.500 rpm separated the solids from the supernatant, thus yielding the crude (native) inulin solution to which sodium azide (0,01%) was added for conservation.

Analysis of the average degree of polymerisation (\overline{DP}) of the inulin was performed by measuring fructose, glucose and saccharose on said supernatant by means of HPLC before and after complete hydrolysis with Fructozym* enzyme (* Trade name of NOVO NORDISK, Denmark) and applying following formula:

$$\frac{Fa - (Fb + GFb/2)}{Ga - (Gb + GFb/2)} + 1 = (\overline{DP}) \text{ inulin}$$

Fa and Fb: fructose respectively after and before hydrolysis

Ga and Gb: glucose respectively after and before hydrolysis

GFb: sucrose before hydrolysis

The value (\overline{DP}) inulin thus obtained corresponds to the degree of polymerisation (\overline{DP}) of the native inulin (which does not take into account the glucose, fructose and saccharose present).

The inulin content was measured by the same method based on the measurement of glucose, fructose and sucrose by HPLC before and after hydrolysis and dividing the result by 1,09 for compensation of the weight gain induced by the hydrolysis. Also the different dilutions, due to the extraction, stabilisation with sodium azide, buffering and enzymatic hydrolysis, were taken into account.

The % inulin (wt % inulin) is obtained by the following formula:

$$\% \text{ inulin} = \left(\frac{Fa - (Fb + GFb/2) + Ga - (Gb + GFb/2)}{1,09} \right) \times \text{dilution factors}$$

The results are given in table 1 below.

5 Table 1: Results on fresh roots:

	harvest date	growing + processing period (days)	inulin content of fresh roots (wt %)	(\overline{DP}) of native inulin
10	17/09	155	17,4	14,2
	01/10	169	17,0	14
	15/10	183	17,2	13,1
	29/10	197	16,3	12,1
	12/11	211	16,5	9,6
15	26/11	225	16,2	8,7
	10/12	231	15,6	8,2

20 B. 300 kg of the roots harvested on 1/10, 29/10 and 26/11 were also processed at pilot scale.

Extraction was done on the sliced chicory roots with a counter current diffuser at 15 kg/hr as used in sugar beet processing (temperature of diffusion: 70-75°C; 125% drawing off; residence time 1hr; pH 5,4-5,8).

25 Depuration was performed by liming and carbonatation similar to beet processing at temperatures between 65 and 75 °C followed by filtration at 85°C.

30 Refining was done by treatment over a strong cation exchanger and a weak anion exchanger (treatment carried out twice), avoiding the pH to drop below 4,5, followed by a treatment with 1% activated carbon and filtered. The resulting syrup was evaporated up to a concentration of 35 Brix, spray dried on a Buchi* (* trade name) lab dryer to particles of 96 % dry matter and analysed. Results are given in table 2 below.

35 More then 80% yield (on inulin basis) of standard grade chicory inulin was thus obtained.

Table 2: Results of standard grade chicory inulin obtained by processing at pilot scale

harvest date	days*	(\overline{DP}) native inulin	(\overline{DP}) standard grade inulin * *	loss in (\overline{DP}) by processing***
1/10	169	14	12,8	1,2
29/10	197	12,1	11,7	1,4
26/11	225	8,7	7,0	1,7

- 10 *total days of growing + processing period; ** (\overline{DP}) after processing
 *** processing as such at pilot scale

C. Manufacture of inulin at industrial scale:

- 15 Roots of chicory consisting mainly of the varieties Madonna, Orchies, and
 Cassel, seeded during April (in average about April 16) in Belgium were
 processed from 19/09 until 15/12 in a factory at Oreya (Belgium) under
 similar conditions as defined above for the process at pilot scale (diffusion at
 72-76 °C; drawing off at 135 %; pH between 5,4-5,8.) Depuration and refining
 20 were done as described above for the process at pilot scale. After treatment
 with active carbon and filtration, processing continued with a filter
 sterilisation step before evaporation. The resulting syrup was spray dried on
 an industrial spray dryer (NIRO * (*Trade name) drying tower), yielding
 particles of 96 % dry matter content, and analysed. The results are shown in
 25 table 3 below. More than 85 % yield (on inulin basis) of standard grade
 inulin was thus obtained.

Table 3: Results of standard grade chicory inulin obtained by processing at industrial scale.

5	harvest date	growing + processing period (days*)	(\overline{DP}) native inulin	(\overline{DP}) standard grade inulin	loss in (\overline{DP}) by processing**
	1/10	169	13,9	12,7	1,2
	15/10	183	12,5	11,2	1,3
10	29/10	197	11,9	10,6	1,3
	12/11	211	9,6	8,1	1,5
	26/11	225	8,9	7,3	1,6
	10/12	231	8,5	6,9	1,6

* calculated from April 16

** processing as such at industrial scale

The mean (\overline{DP}) of the standard grade chicory inulin manufactured at industrial scale (mean value of (\overline{DP}) taken over the indicated period) thus obtained is shown in table 4.

Table 4: Mean (\overline{DP}) of standard grade chicory inulin

25	period taken	days of processing period	mean (\overline{DP}) native inulin	mean (\overline{DP}) standard grade inulin
	1/10->12/11	43	12,0	10,7
	1/10->26/11	57	11,4	10,0
	1/10->10/12	71	10,9	9,5

It is general practice in Belgium and in most parts of the northern hemisphere to stop the production of inulin before the end of November in order to be able to sell standard grade chicory inulin with a (\overline{DP}) of about 10. This requirement of (\overline{DP}) would not be reached including product from later harvesting dates.

Example 2:

Chicory inulin from chicory roots grown in California (USA) under climatological temperature conditions according to the invention (fall harvest)

5

Chicory, cultivar Orchies, coated seeds, were planted on 26/03 in Dixon (USA) under irrigated conditions. Stand was 85.000 roots/ha. Harvest was done manually, about monthly from 21/8 until 26/11 and the roots were processed without delay. Analysis of the native inulin was done according to the method described in Example 1. The results are shown in table 5.

10

Table 5 : Results on fresh roots:

15	harvest	growing +	inulin content	(\overline{DP})
	date	processing period (days)	of fresh roots (wt %)	native inulin
	21/8	149	15.8	16.6
	24/9	183	17.1	18.0
	29/10	218	16.5	17.9
20	26/11	246	15.7	17.5

Example 3:

Chicory inulin from chicory roots grown in California (USA) under climatological temperature conditions according to the invention (fall harvest)

25

Chicory, cultivar Cassel, coated seeds, were planted on 04/04 in Artois (USA) under irrigated conditions. Harvest was done manually, about monthly from 21/08 until 26/11, and the roots were processed without delay. Analysis of the native inulin was done according to the method indicated in example 1. The results are shown in table 6 below.

30

Table 6: Results on fresh roots:

	harvest date	growing + processing period (days)	inulin content of fresh roots wt %	(\overline{DP}) native inulin
5				
	21/08	140	15,3	15,0
	19/09	169	16,7	17,4
	24/10	204	17,0	18,2
10	26/11	237	17,1	15,4

Example 4:

Chicory inulin from chicory roots grown in California (USA) under climatological temperature conditions according to the invention (Spring harvest)

- 15 a) Chicory, cultivar Orchies, coated seeds were planted on 3/10 and 30/10 in Davis (California) under irrigated conditions. Harvest was done manually begin May and June of the following year, and the roots were analysed without delay, according to the method indicated in example 1. The results are shown in table 7 below.

Table 7: Results on cultivar Orchies (Spring harvest in Davis, USA)

	planting date	harvest date	growing + processing period (days)	inulin content of fresh roots wt %	(\overline{DP}) native inulin
25					
	3/10	8/05	217	13,7	16,0
30		2/06	242	14,5	16,0
	30/10	3/06	216	13,5	14,6

- 35 b) Chicory, cultivar Orchies and Madonna , coated seeds were planted on 2/10 and 31/10 in Coalinga (California) under irrigated conditions. Harvest was done manually begin and end of June of the following year, and the roots were analysed without delay, according to the method indicated in example 1. The results are shown in table 8 below.

Table 8: Results on cultivars Orchies and Madonna
(Spring harvest in Coalinga, USA)

	planting date	harvest date	growing + processing period (days)	inulin content of fresh roots (wt %)	(\overline{DP}) native inulin
5	ORCHIES:				
	2/10	9/06	250	13,7	17,1
10		29/06	270	16,9	19,6
	31/10	9/06	221	14,9	19
		29/06	241	17,0	17,5
	MADONNA:				
15	2/10	9/06	250	14,2	18,1
		29/06	270	14,7	20,6
	31/10	9/06	221	13,0	16,8
		29/06	241	13,8	18,8

20 Example 5:

Production of depurated inulin from chicory roots grown in Artois
(California; USA) under climatological temperature conditions according to
the invention (Fall harvest)

25 15 kg chicory roots from Example 3 (Cassel cultivar; harvest on 26/11; with
 inulin content 17,1 % and (\overline{DP}) native inulin 15,4) were processed according
 to Example 1 into a crude (native) inulin syrup: the extraction was done on
 3 batches of 5 kg roots which were sliced into small pieces of 0,3-0,8 cm thick,
 0,3-0,8 cm wide and 2 to 6 cm long. The slicings were extracted by treatment
 30 with 7 kg boiling water and keeping the mixture at 75°C for 90 minutes at
 pH 5,4-5,8. The slicings were separated from the extract and the turbid liquid
 so obtained from the 3 batches were combined for further treatment.

In total 18,1 kg of extraction juice were obtained, representing about 2.200
 grams of crude inulin. This juice could be stored until further processing by
 35 adding 0,01 % sodium azide.

Processing was continued by a depuration step comprising :

- heating of the juice to 85 °C for 15 minutes for resolubilisation of the inulin,
- cooling of the juice to 65°C and prelining,

- a first liming - carbonatation at 70-75°C, with filtration at 80-85°C, followed by a second liming - carbonatation at 70-75°C with filtration at 80-85 °C. The solution obtained was then concentrated by evaporation from about 15 Brix to about 32 Brix before further processing. In total 7,4 kg depurated juice at 32 Brix were obtained showing a 85 % inulin purity. If processing was not continued without delay, the juice was kept frozen till further processing.

Example 6:

Manufacture of standard grade inulin from chicory roots grown in Artois (California: USA) under climatological temperature conditions according to the invention (Fall harvest)

3 kg of depurated juice at 32 Brix obtained in example 5 was heated at 95 °C for 20 minutes, cooled down to 35 °C and sent subsequently over 1 l of strong cation exchanger, 1 l of weak anion exchanger, 0,5 l strong cation exchanger and 0,5 l of weak anion exchanger. Accordingly, the solution was treated with 1% powdered activated carbon at 60°C, filtered, evaporated to 35 Brix and spray dried in a laboratory Buchi* spray drier (* Trade name) yielding particles with a 96 % dry matter content.

The obtained product is improved standard grade chicory inulin with a purity of more then 99,7 % inulin and a (\overline{DP}) of 14,2; (the loss of (\overline{DP}) due to processing was thus 1,2). The results of the analysis of the product performed by HPLC are given in table 9 below.

Table 9: Composition of Improved standard grade chicory inulin
(Results in % of total dry substance)

	<u>Improved standard grade inulin</u>	<u>standard grade inulin *</u>
	%	%
Fructose	1,1	3,4
Glucose	0,5	1,2
Sucrose	4,7	10,2
Inulin	93,3	85
Total	99,6	99,8

* comparative data obtained by applying the same procedure to crude inulin solution from example 1, harvest 29/10.

Extrapolating these results to Example 3 over a processing period (including harvesting and processing) from 19/9 to 26/11 results in a mean (\overline{DP}), thus taken over a processing period of 69 days, for the improved standard grade chicory inulin of 16,5.

5

Example 7:

Manufacture of high performance grade inulin from chicory roots grown in Artois (California; USA) under climatological temperature conditions according to the invention (Fall harvest)

10 The method used is according to the one described in patent application WO 96/01849, enclosed herein by reference, involving a directed crystallisation of inulin from a meta-stable aqueous polydisperse inulin solution.

1 kg of depurated juice at 32 Brix obtained in example 5 , containing 270 grams of depurated inulin was poured in a Duran* (* Trade name) bottle and heated for 15 min in a pressure cooker to completely solubilise the inulin. Accordingly, this solution is cooled down to 20°C within a period of 5 minutes directly followed by adding an inulin grafting suspension in a ratio of 1/500 wt% (grafting dry matter/inulin dry matter). The solution was left for 20 hours for crystallisation, then filtered on a vacuum filter and the filter cake was washed with 330 grams of distilled water, yielding a filter cake at 42 % dry matter. After drying 127 grams of high performance (HP) grade chicory inulin, with a purity of >99,5% and a (\overline{DP}) of 29, were obtained. This corresponds to a yield of 47 % (based on inulin).

When the same procedure (depuration and directed crystallisation) is applied to a crude inulin solution (from example 1; harvest 29/10) resulting from roots with a (\overline{DP}) native inulin of 12,1, then the resulting high performance grade inulin has a (\overline{DP}) of 25 and is obtained in a yield of only 29% (based on inulin).

30 Example 8:

Manufacture of fructo-oligosaccharides from chicory roots grown in Artois (California; USA) under climatological temperature conditions according to the invention (Fall harvest)

1 kg of depurated juice at 32 Brix obtained in example 5 was heated at 95 °C for 20 minutes, cooled down to 60°C, pH adjusted to 5,4 with HCl and 160 Units of a preparation of an endo-inulinase (obtained by fractionation of an

35

inulinase of *Aspergillus ficuum* as described by B. Norman et al., Denpun Kagaku, Vol., 36, 103-111, (1989)) was added according to the process described in patent application PCT/BE97/00087, enclosed herein by reference.

- 5 The juice was incubated then for 24 hr at 60°C, boiled for 2 minutes to deactivate the enzymes, cooled to 15°C and further refined as described in example 1, yielding an improved fructo-oligosaccharide (oligofructose) composition. The results of the analysis, performed by gaschromatography, of the obtained composition, in comparison to the composition obtained
10 from crude inulin juice (from example 1) coming from roots with a (\overline{DP}) native inulin of 12,1, are given below in Table 10.

Table 10: Composition of improved Fructo-oligosaccharide

(Results in % of total dry substance (measured as Brix))

	Improved oligofructose composition	standard oligofructose composition *
15	Fructose 4,0	4,4
	Glucose 0,5	1,0
	Sucrose 4,2	9,4
20	other DP2 3,3	2,8
	GF2=DP3 2,8	6,3
	F3=DP3 27,5	21,8
	GF3=DP4 7,5	11,4
	F4=DP4 20,1	13,4
25	GF4=DP5 10,0	12,4
	F5=DP5 6,5	3,8
	GF5=DP6 7,8	9,4
	F6=DP6 3,8	1,9
	GF6=DP7 1,1	1,3
30	F7=DP7 0,4	0
	GF7=DP8 0,3	0,3
	GF8=DP9 0,1	0,1
	GF9=DP10 0,1	0,0
	TOTAL 99,9	99,7
35	% OLIGOFRUCTOSE 91,2	84,9

* comparative data obtained by applying the same procedure to crude inulin solution from example 1; harvest 29/10.

Example 9:

Manufacture of fructose from chicory roots grown in Artois (California: USA) under climatological temperature conditions according to the invention (Fall harvest)

5

1 kg of depurated juice at 32 Brix obtained in example 5 was heated at 95 °C for 20 minutes, cooled down to 60°C, pH adjusted to 4,5 with HCl and 775 Units of Fructozym* enzyme (Trade name; NOVO NORDISK, Denmark) was added . The mixture was incubated for 24 hr at 60°C, boiled for 2 minutes to deactivate the enzymes, cooled to 15°C and further refined as described in example 1 yielding an improved fructose composition. The results of the analysis, performed by HPLC, of the obtained composition, in comparison to the composition obtained from crude inulin juice (from example 1) coming from roots with a (\overline{DP}) native inulin of 12,1, are given below in Table 11.

15

Table 11: Composition of improved Fructose

(Results in % of total dry substance (measured as Brix))

20

	Improved fructose composition	standard fructose composition*
Fructose	90,0	86,5
Glucose	8,2	11,3
Sucrose	0,3	0,4
25 Difructosedianhydride	1,3	1,6
DP3	0	0
Total	99,8	99,8

* comparative data

30

CLAIMS

1. Process for the manufacture of chicory inulin from chicory roots by conventional manufacturing techniques, wherein the source material for the process are roots of chicory which have been grown in appropriate regions and grown and processed under proper climatological temperature conditions which are such that during a period of at least from the beginning of the third month of the growing period till the end of the processing of the chicory roots the FEH gene in the chicory roots has not been triggered by the occurrence of low temperature conditions, said chicory roots have had a growing period of at least 150 days, and said chicory has been seeded in the northern hemisphere within a period selected from the periods ranging from December 1 till March 14, from May 15 till May 31, from June 1 till June 14, and from June 15 till November 30, provided that when said chicory has been seeded in the periods from May 15 till May 31, and from June 1 till June 14, the chicory roots have had a growing period of at least 180 days, or in the southern hemisphere within a period selected from the periods ranging from June 1 till September 14, from September 15 till September 30, from October 1 till November 14, from November 15 till November 30, and from December 1 till May 31.

2. Process according to claim 1, wherein the chicory has had a growing period of at least 180 days.

3. Process according to claim 1 or claim 2 wherein the chicory inulin is standard grade or low sugar standard grade chicory inulin with a (\overline{DP}) of at least 10, improved standard grade or improved low sugar standard grade chicory inulin with a mean (\overline{DP}) of at least 12, high performance grade chicory inulin with a (\overline{DP}) of at least 20, or improved high performance grade chicory inulin with a mean (\overline{DP}) of at least 20, the mean (\overline{DP}) being taken over a processing period of at least 60 days.

4. Process for the manufacture of improved standard grade chicory inulin from chicory roots through conventional manufacturing techniques, wherein the source material for the process are roots of chicory grown in appropriate regions and which have been grown and processed under proper climatological temperature conditions wherein, within a period of at least 220 consecutive days immediately preceding the end of the processing of the roots, no low temperature conditions occurred which triggered the

FEH gene in chicory roots, said roots have had a growing period of at least 160 days, the FEH gene in the chicory roots has not been triggered by the occurrence of low temperature conditions within the period from at least the beginning of the third month of the growing period till the end of the processing of the roots, and the product obtained is improved standard grade chicory inulin with a mean (\overline{DP}) taken over a processing period of at least 60 days, which is at least 12.

5 5. Process according to claim 4, wherein the climatological temperature conditions are such that no low temperature conditions which trigger the FEH gene in chicory roots occur within a total period of at least 240 consecutive days, the chicory has had a growing period of at least 180 days, and the product obtained is improved standard grade chicory inulin with a mean (\overline{DP}), taken over a processing period of at least 60 days, which is at least 12.

10 6. Process according to claim 4 or claim 5 wherein the chicory has been seeded in the northern hemisphere within a period selected from the periods ranging from December 1 till March 14, from March 15 till May 14, from May 15 till May 31, and from June 1 till November 30, or in the southern hemisphere within a period selected from the periods ranging from June 1 till September 14, from September 15 till November 14, from November 15 till November 30, and from December 1 till May 31.

15 7. Process according to any of claims 1 to 6 wherein the said proper climatological temperature conditions are such that during the concerned period immediately preceding the end of the processing of the chicory roots the temperature in the temperature shelter has not dropped below minus 1°C.

20 8. Process according to any of claims 1 to 7 wherein said appropriate regions for growing chicory which present, besides the known conditions for growing chicory, said proper climatological temperature conditions, comprise a region selected from the group consisting of the Californian region of the USA, certain valleys of Chile, and certain maritime regions of Europe, Turkey, Australia, Morocco, Algeria, South Africa, China and India.

25 9. Process according to any of claims 1 to 8 for the manufacture of standard grade chicory inulin or improved standard grade chicory inulin, wherein the conventional manufacturing techniques comprise the following steps:

(i) isolation of the inulin from the chicory roots yielding an aqueous solution of crude inulin,

(ii) purification of the crude inulin obtained in step (i) yielding an aqueous solution of purified inulin, optionally followed by concentration of this solution by partial removal of the water yielding a purified inulin concentrate, and

(iii) isolation in particulate form of the inulin from the aqueous solution or concentrate of purified inulin obtained in step (ii), thereby yielding, respectively, standard grade chicory inulin or improved standard grade chicory inulin.

10. Process according to claim 9, wherein the manufacturing techniques comprise:

- for step (i): extraction with hot water of the inulin from fresh slices or shreds of the chicory roots, yielding an aqueous solution of crude inulin,

- for step (ii): purification of the aqueous solution of crude inulin obtained in step (i) by depuration followed by refining, and

- for step (iii): isolation of, respectively, standard grade chicory inulin or improved standard grade chicory inulin, in particulate form by spray drying.

11. Process for the manufacture of low sugar standard grade chicory inulin or improved low sugar standard grade chicory inulin containing in total less than 1 weight % monomeric saccharides and sucrose, by conventional techniques from chicory roots, wherein the source material are chicory roots which have been grown and processed under the conditions as defined in any of claims 1 to 8.

12. Process for the manufacture of low sugar standard grade chicory inulin or improved low sugar standard grade chicory inulin containing in total less than 1 weight % monomeric saccharides and sucrose, wherein, respectively, standard grade chicory inulin or improved standard grade chicory inulin or the corresponding intermediate purified inulin, obtained by a process defined in any of claims 1 to 10 is used as a source material and is, in accordance with known techniques, subjected to the following additional consecutive steps:

(iv) removal of the monomeric saccharides and sucrose, yielding a low sugar inulin solution or concentrate, and

(v) isolating the low sugar inulin in particulate form from the solution or concentrate obtained in step (iv),

and the product obtained is, respectively, low sugar standard grade chicory inulin with a mean (\overline{DP}) of at least 10 or improved low sugar standard grade chicory inulin with a mean (\overline{DP}) of at least 12, the mean (\overline{DP}) being taken over a processing period of at least 60 days.

5 13. Process for the manufacture by conventional techniques from chicory roots of high performance grade chicory inulin with a (\overline{DP}) of at least 20, or improved high performance grade chicory inulin with a mean (\overline{DP}), taken over a processing period of the chicory roots of at least 60 days, of at least 20, which are essentially free from low molecular monomeric
10 saccharides, dimeric saccharides and oligofructose, and essentially free from colourings, salts, proteins, organic acids and technological aids, wherein the source material are chicory roots which have been grown and processed under the conditions as defined in any of claims 1 to 2, and 4 to 8.

15 14. Process for the manufacture of high performance grade chicory inulin with a (\overline{DP}) of at least 20, or of improved high performance grade chicory inulin with a mean (\overline{DP}), taken over a processing period of the chicory roots of at least 60 days, of at least 20, which are essentially free from low molecular monomeric saccharides, dimeric saccharides and
20 oligofructose, and essentially free from colourings, salts, proteins, organic acids and technological aids, wherein standard grade chicory inulin with a (\overline{DP}) of at least 12, respectively improved standard grade chicory inulin with a mean (\overline{DP}), taken over a processing period of the chicory roots of at least 60 days, of at least 12, or its intermediate, depurated or refined inulin, obtained by a process defined in any of claims 1 to 10, is used as a source
25 material and subjected, in accordance with known techniques, to the following consecutive steps:

(vi) fractionation, and

(vii) isolation in particulate form of the high performance grade inulin from the fractionated product obtained in step (vi),
30 thereby providing high performance grade chicory inulin, respectively improved high performance grade chicory inulin in a yield of at least 40 % based on the source inulin.

15 15. Process according to claim 14, wherein the fractionation is carried out by directed crystallisation of an aqueous metastable solution of the source material, and the isolation of the fractionated inulin in particulate
35

form is carried out by filtration or centrifuging including washing with water.

16. Process according to any of claims 13 to 15, wherein the source inulin has a (\overline{DP}), respectively a mean (\overline{DP}) of at least 14, and the high performance grade chicory inulin, respectively improved high performance grade chicory inulin, is obtained in a yield of at least 45 % based on the source material, and has a (\overline{DP}), respectively a mean (\overline{DP}), of at least 20, the mean (\overline{DP}) being taken over a processing period of the source chicory roots of at least 60 days.

17. Process for the manufacture of a partial hydrolysate of chicory inulin, by conventional techniques from chicory roots, wherein the source material are chicory roots which have been grown and processed under the conditions as defined in any of claims 1 to 2, and 4 to 8 and the product obtained is a polydisperse oligofructose composition.

18. Process for the manufacture of a partial hydrolysate of chicory inulin by conventional techniques, wherein respectively, standard grade or improved standard grade chicory inulin or the corresponding intermediates, depurated or refined inulin, obtained by a process defined in any of claims 1 to 10, is used as a source material, and the product obtained is a polydisperse oligofructose composition containing at least 90 % by weight dry substance and the oligofructose has a (DP) from 2 to 10.

19. Process for the manufacture of a complete hydrolysate of chicory inulin, by conventional techniques from chicory roots, wherein the source material are chicory roots which have been grown and processed under the conditions as defined in any of claims 1 to 2, and 4 to 8, and the product obtained is a fructose composition.

20. Process for the manufacture by known techniques of a complete hydrolysate of inulin, wherein, respectively, standard grade or improved standard grade inulin or the corresponding intermediate, depurated or refined inulin, obtained by a process defined in any of claims 1 to 10, is used as a source material, and the product obtained is a fructose composition containing at least 89 % by weight fructose, calculated on dry substance.

21. Process for the manufacture of a derivative of chicory inulin, by conventional techniques from chicory inulin or an intermediate thereof, wherein the source material for the inulin are chicory roots which have

been grown and processed under the conditions as defined in any of claims 1 to 2, and 4 to 8.

22. Improved standard grade chicory inulin with an inulin content, expressed on dry matter, of at least 92% by weight, and a maximal total content of glucose, fructose and sucrose of 8 % by weight, characterised in that the inulin has a mean (\overline{DP}) taken over a processing period of the source chicory roots of at least 60 days, of at least 12, obtainable by a process according to any of claims 1 to 10.

23. Improved low sugar standard grade chicory inulin with, expressed on dry matter, an inulin content of more than 99% by weight, and a total content of glucose, fructose and sucrose of less than 1% by weight, characterised in that the inulin has a mean (\overline{DP}) taken over a processing period of the source chicory roots of at least 60 days, of at least 12, obtainable by a process according to any of claims 1 to 3 and 11 to 12.

24. Improved high performance grade chicory inulin which is essentially free from low molecular monomeric saccharides, dimeric saccharides and oligofructose, and essentially free from colourings, salts, proteins, organic acids and technological aids, with a mean (\overline{DP}), taken over a processing period of the source chicory roots of at least 60 days, of at least 20, obtainable by a process according to any of claims 1 to 3 and 13 to 16.

25. Improved polydisperse oligofructose composition containing at least 90 % by weight dry substance and wherein the oligofructose has a (DP) from 2 to 10, obtainable by a process according to claim 17 or 18.

26. Improved fructose composition containing at least 89 % by weight of fructose and having a ratio fructose/glucose of at least 90/10, obtainable by a process according to claim 19 or 20.

27. Method of use of roots of chicory as a source material in a process for the manufacture of chicory inulin, of a partial or a complete hydrolysate of inulin, or a derivative of inulin, characterised in that the roots have been grown in appropriate regions and grown and processed under proper climatological temperature conditions as defined in any of claims 1 to 2, with optionally a processing period of said roots of at least 60 days adjacent to the growing period, as well as under the conditions defined in any of claims 4 to 8.

28. Method of use of improved standard grade chicory inulin, improved low sugar standard grade chicory inulin, improved high

performance grade chicory inulin, an improved polydisperse oligofructose composition or an improved fructose composition, as defined in any of claims 22 to 26, as an ingredient in the manufacture of food, feed, drinks, a prophylactic or a therapeutic composition, or a non-food composition.



ABSTRACT

5

The invention provides an improved process for the manufacture of chicory inulin, hydrolysates and derivatives of inulin by conventional manufacturing techniques from roots of chicory grown in appropriate regions and which have been grown and processed under proper climatological temperature conditions. Selection of said proper conditions enables to provide a growing and/or processing period for the chicory roots which may partly or wholly extend beyond the conventional periods.

10

15

The invention relates also to improved grades of chicory inulin, of hydrolysates of chicory inulin, such as e.g. a polydisperse oligofructose composition and a fructose composition, as well as to the use of these products in the manufacture of food, feed, drinks, prophylactic and therapeutical compositions, chemical derivatives and non-food compositions

20

Improved standard grade chicory inulin according to the invention presents a degree of polymerisation (\overline{DP}) which is at least 20 % higher than the one of conventional standard grade chicory inulin.

performance grade chicory inulin, an improved polydisperse oligofructose composition or an improved fructose composition, as defined in any of claims 22 to 26, as an ingredient in the manufacture of food, feed, drinks, a prophylactic or a therapeutic composition, or a non-food composition.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/00155

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C08B37/18 A01G1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01G C08B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MICHAEL ERNST ET AL.: "Carbohydrate changes in chicory during growth and storage." SCIENTIA HORTICULTURAE, vol. 63, 1995, pages 251-261, XP002069326 see page 252, paragraph 2.1 see page 259 --- -/--	1-28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 May 1999

Date of mailing of the international search report

07/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lensen, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/00155

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
Y	CHEMICAL ABSTRACTS, vol. 127, no. 18, 3 November 1997 Columbus, Ohio, US; abstract no. 245507, BAERT JOOST R.A.: "The effect of sowing and harvest date and cultivar on inulin yield and composition of chicory roots." XP002069328 see abstract & IND. CROPS PROD., vol. 6, no. 3,4, 1997, pages 195-199, ---	1-28
Y	WIM VAN DEN ENDE ET AL.: "Fructan synthesizing and degrading activities in chicory roots (Cichorium intybus L.) during field-growth, storage and forcing." PLANT PHYSIOL., vol. 149, 1996, pages 43-50, XP002069327 STUTTGART see summary ---	1-28
Y	EP 0 787 745 A (MCNEIL-PPC) 6 August 1997 see page 9, line 15 - line 32 see page 2, line 38 - line 45 ---	1-28
A	LEEN DE LEENHEER ET AL.: "Progress in the elucidation of the composition of chicory inulin." STARCH/ DIE STÄRKE, vol. 46, no. 5, 1 May 1994, pages 193-196, XP000442587 WEINHEIM, DE -----	22-28

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/00155

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 787745	A	06-08-1997	
		AU 1249597 A	14-08-1997
		BR 9700887 A	27-10-1998
		CA 2196739 A	06-08-1997
		HU 9700359 A	30-03-1998
		JP 9324002 A	16-12-1997
		NO 970490 A	06-08-1997
		NZ 314187 A	28-01-1999

